

ADAS Subroutine bxstkc

```
      SUBROUTINE BXSTKC ( NDLEV , NDMET ,  
&                        NORD   , NMET   ,  
&                        IORDR  , IMETR  ,  
&                        CC     , STCK   ,  
&                        CRED  
&                        )
```

```
C-----  
C  
C *****  
C ***** FORTRAN77 SUBROUTINE: BXSTKC *****  
C *****  
C  
C PURPOSE: TO STACK UP IN 'CRED' THE TRANSITION RATE BETWEEN METASTA-  
C           BLE LEVELS FOR A GIVEN TEMPERATURE STABLE LEVEL FOR A GIVEN  
C           TEMPERATURE AND DENSITY.  
C  
C CALLING PROGRAM:  ADAS205/ADAS206  
C  
C SUBROUTINE:  
C  
C INPUT : (I*4)  NDLEV  = MAXIMUM NUMBER OF ENERGY LEVELS ALLOWED  
C INPUT : (I*4)  NDMET  = MAXIMUM NUMBER OF METASTABLE LEVELS ALLOWED  
C  
C INPUT : (I*4)  NORD   = NUMBER OF ORDINARY EXCITED LEVELS  
C INPUT : (I*4)  NMET   = NUMBER OF METASTABLE LEVELS  
C  
C INPUT : (I*4)  IMETR() = INDEX OF METASTABLE IN COMPLETE LEVEL LIST  
C                       (ARRAY SIZE = 'NDMET' )  
C INPUT : (I*4)  IORDR() =INDEX OF ORDINARY EXCITED LEVELS IN COMPLETE  
C                       LEVEL LIST.  
C                       (ARRAY SIZE = 'NDLEV' )  
C  
C INPUT : (R*8)  CC(, )  = RATE MATRIX COVERING ALL TRANSITIONS  
C                       (UNITS: SEC-1)  
C                       VALUES FOR GIVEN TEMPERATURE AND DENSITY.  
C                       1st DIMENSION: ENERGY LEVEL INDEX  
C                       2nd DIMENSION: ENERGY LEVEL INDEX  
C INPUT : (R*4)  STCK(, ) = POPULATION MATRIX COVERING ALL NON-METAST-  
C                       ABLE/ORDINARY EXCITED LEVELS AS FUNCTION  
C                       OF METASTABLE INDEX.  
C                       VALUES FOR GIVEN TEMPERATURE AND DENSITY.  
C                       1st DIMENSION: ORDINARY EXCITED LEVEL INDEX  
C                       2nd DIMENSION: METASTABLE LEVEL INDEX  
C  
C OUTPUT: (R*8)  CRED(, ) = MATRIX OF TRANSITION RATES BETWEEN  
C                       METASTABLE LEVELS.  
C                       (UNITS: SEC-1)  
C                       VALUES FOR GIVEN TEMPERATURE AND DENSITY.  
C                       1st DIMENSION: METASTABLE LEVEL INDEX  
C                       2nd DIMENSION: METASTABLE LEVEL INDEX  
C  
C  
C           (I*4)  IM1    = METASTABLE LEVEL ARRAY INDEX  
C           (I*4)  IM2    = METASTABLE LEVEL ARRAY INDEX  
C           (I*4)  IS     = ORDINARY EXCITED LEVEL INDEX
```

```

C
C
C ROUTINES: NONE
C
C NOTE:
C      CRED(IM1,IM2) = ( the transition rate from IM2 to IM1 )
C                          +
C                          SUM( (the transistion rate from ordinary
C                              level IS to IM1) x (the population
C                              in metastable level IM2 that excite
C                              to oridinary level IS) )
C
C                          ABOVE SUM IS OVER ALL ORDINARY LEVELS.
C
C
C AUTHOR:  PAUL E. BRIDEN (TESSELLA SUPPORT SERVICES PLC)
C          K1/0/81
C          JET EXT. 4569
C
C DATE:    09/10/90
C
C UPDATE:  20/05/93 - P BRIDEN: STCK ARRAY CHANGED FROM REAL*8 -> REAL*4
C
C-----
C
C-----
C
C          INTEGER          IMETR (NDMET) ,          IORDR (NDLEV)
C          INTEGER          NDLEV,          NDMET,          NMET,          NORD
C          REAL*8           CC (NDLEV, NDLEV) ,          CRED (NDMET, NDMET)
C          REAL             STCK (NDLEV, NDMET)

```